


Library
National Wetlands Research Center
U. S. Fish and Wildlife Service
700 Cajundome Boulevard
Lafayette, La. 70506

FWS/OBS-82/10.84
SEPTEMBER 1984

HABITAT SUITABILITY INDEX MODELS: LAKE TROUT (EXCLUSIVE OF THE GREAT LAKES)



Fish and Wildlife Service

Department of the Interior

SK
361
.U54
no.82-
10.84

MODEL EVALUATION FORM

Habitat models are designed for a wide variety of planning applications where habitat information is an important consideration in the decision process. However, it is impossible to develop a model that performs equally well in all situations. Assistance from users and researchers is an important part of the model improvement process. Each model is published individually to facilitate updating and reprinting as new information becomes available. User feedback on model performance will assist in improving habitat models for future applications. Please complete this form following application or review of the model. Feel free to include additional information that may be of use to either a model developer or model user. We also would appreciate information on model testing, modification, and application, as well as copies of modified models or test results. Please return this form to:

Habitat Evaluation Procedures Group
U.S. Fish and Wildlife Service
2627 Redwing Road, Creekside One
Fort Collins, CO 80526-2899

Thank you for your assistance.

Species _____ Geographic Location _____

Habitat or Cover Type(s) _____

Type of Application: Impact Analysis ____ Management Action Analysis ____
Baseline ____ Other _____

Variables Measured or Evaluated _____

Was the species information useful and accurate? Yes ____ No ____

If not, what corrections or improvements are needed? _____

Were the variables and curves clearly defined and useful? Yes No

If not, how were or could they be improved? _____

Were the techniques suggested for collection of field data:

Appropriate? Yes No

Clearly defined? Yes No

Easily applied? Yes No

If not, what other data collection techniques are needed? _____

Were the model equations logical? Yes No

Appropriate? Yes No

How were or could they be improved? _____

Other suggestions for modification or improvement (attach curves, equations, graphs, or other appropriate information) _____

Additional references or information that should be included in the model:

Model Evaluator or Reviewer _____ Date _____

Agency _____

Address _____

Telephone Number Comm: _____ FTS _____

FWS/OBS-82/10.84
September 1984

HABITAT SUITABILITY INDEX MODELS: LAKE TROUT
(EXCLUSIVE OF THE GREAT LAKES)

by

Michael D. Marcus
Western Aquatics, Inc.
P.O. Box 546
Laramie, WY 82070

Wayne A. Hubert
and
Stanley H. Anderson
Wyoming Cooperative Fishery and
Wildlife Research Unit
University of Wyoming
Laramie, WY 82071

Performed for

Western Energy and Land Use Team
Division of Biological Services
Research and Development
Fish and Wildlife Service
U.S. Department of the Interior
Washington, DC 20240

This report should be cited as:

Marcus, M. D., W. A. Hubert, and S. H. Anderson. 1984. Habitat suitability index models: Lake trout (Exclusive of the Great Lakes). U.S. Fish Wildl. Serv. FWS/OBS-82/10.84. 12 pp.

PREFACE

The Habitat Suitability Index (HSI) models presented in this publication aid in identifying important habitat variables. Facts, ideas, and concepts obtained from the research literature and expert reviews are synthesized into HSI models, which are scaled to produce an index between 0 (unsuitable habitat) and 1 (optimal habitat). The models are hypotheses of species-habitat relationships, and model users should recognize that the degree of veracity of the HSI model and assumptions will vary according to geographical area and the extent of the data base for individual variables. After clear study objectives have been set, the HSI model building techniques presented in U.S. Fish and Wildlife Service (1981)¹ and the general guidelines for modifying HSI models and estimating model variables presented in Terrell et al. (1982)² may be useful for simplifying and applying the models to specific impact assessment problems. Simplified models should be tested with independent data sets, if possible. A statistically-derived model that is an alternative to using Suitability Indices to calculate an HSI is referenced in the text.

Model reliability is likely to vary in different geographical areas and situations. The U.S. Fish and Wildlife Service encourages model users to provide comments, suggestions, and test results that may help us increase the utility and effectiveness of this habitat-based approach to impact assessment. Please send comments to:

Habitat Evaluation Procedures Group
Western Energy and Land Use Team
U.S. Fish and Wildlife Service
2627 Redwing Road
Ft. Collins, CO 80526-2899

¹U.S. Fish and Wildlife Service. 1981. Standards for the development of habitat suitability index models. 103 ESM. U.S. Fish Wildl. Serv., Div. Ecol. Serv. n.p.

²Terrell, J. W., T. E. McMahon, P. D. Inskip, R. F. Raleigh, and K. L. Williamson (1982). Habitat suitability index models: Appendix A. Guidelines for riverine and lacustrine applications of fish HSI models with the Habitat Evaluation Procedures. U.S. Fish Wildl. Serv. FWS/OBS-82/10.A. 54 pp.



CONTENTS

	<u>Page</u>
PREFACE	iii
ACKNOWLEDGMENTS	vi
HABITAT USE INFORMATION	1
General	1
Age, Growth, and Food	2
Reproduction	3
Specific Habitat Requirements	4
HABITAT SUITABILITY INDEX (HSI) MODELS	5
Model Applicability	5
Model Description	5
Suitability Index (SI) Graphs for Model Variables	5
Development of Suitability Index Graphs: Rationale and Assumptions	7
HSI Model for Stocked Lake Trout Populations	9
HSI Model for Reproducing Lake Trout Populations in Lakes	9
HSI Model for Reproducing Lake Trout Populations in Reservoirs	9
Interpreting Model Outputs	9
ADDITIONAL HABITAT MODEL	10
REFERENCES	10

ACKNOWLEDGMENTS

We wish to thank members of the Western Energy and Land Use Team for helpful advice on model construction, as well as Thomas Edsall, Steven Grabowski, Thomas Pettengill, John Keifling, and Eric Bergersen for reviewing the model. Word processing was provided by Carolyn Gulzow and Dora Ibarra. Editing was provided by Cathy Short and Tricia Rosenthal. The cover of this document was illustrated by Jennifer Shoemaker.

LAKE TROUT (Salvelinus namaycush)

HABITAT USE INFORMATION

General

The lake trout is an important commercial and sport fish in North America. In the Central Rocky Mountain region, lake trout are commonly referred to as "mackinaw". There is good evidence that lake trout should be called "lake charr" (Morton 1980). No subspecies of lake trout is presently recognized (Robins et al. 1980). The species, however, has extreme variability throughout its range, making it difficult to draw general conclusions about its biology (Martin and Olver 1980).

Native lake trout are confined to northern North America, with their distribution in the United States limited to the Great Lakes drainage and parts of northern New England, New York, Wisconsin, Minnesota, Montana, and Alaska (Carlander 1969; Martin and Olver 1980). In Canada, native lake trout occur in all provinces and territories, except Prince Edward Island, insular Newfoundland, portions of the prairie provinces, and coastal British Columbia. Lake trout have been introduced into California, Nevada, Colorado, Washington, Oregon, Idaho, Utah, Wyoming, Massachusetts, Connecticut, Argentina, Peru, Bolivia, Finland, Sweden, Switzerland, France, and New Zealand (Martin and Olver 1980). For the most part, lake trout are confined to fresh water, though they may enter brackish water in the far northern part of their range.

Morphometric and meristic data indicate that the distribution of native lake trout resulted from migrations originating from at least three and likely five unglaciated refugia following the Wisconsin Glaciation (Khan and Qadri 1971; Martin and Olver 1980). Genetic differences present in these various historical population stocks may partially account for the great biological variability observed in this species.

Lake trout are highly mobile, with individuals making horizontal and vertical movements in response to various physical, chemical, and biological factors (Martin and Olver 1980). While few data exist on the movement of young lake trout, it appears that fry move from the spawning areas within a month after hatching and travel to the deeper waters of the lake. This may be to avoid higher light intensities or predation (Martin and Olver 1980). Sometime during the first 22 weeks after hatch, the alevins fill their swim-bladders with a trip to the surface. Young-of-the-year and older lake trout tend to remain in the deeper waters of lakes.

Individual lake trout may occur wherever water temperatures are favorable. For example, Johnson (1975) reported that the youngest trout in Great Bear Lake used a variety of habitats, including rocky shore lines, inflowing streams, and deep waters. In inland lakes, the vertical movement of lake trout appeared to be dependent primarily upon water temperature (Martin and Olver 1980). Dahlberg (1981) reported that temperature was the most important factor controlling the depth distribution of lake trout in Lake Cayuga, New York.

While lake trout tend to be nomadic, some individuals, particularly those from planted stock, appear sedentary (Martin and Olver 1980). For comparison, one individual tagged in Lake Superior travelled 306 km in 19 days (Eschmeyer et al. 1953). In contrast, other lake trout planted in Lake Superior were found after 3 years to be most abundant at distances of 3.2 to 6.4 km from the planting area (Pycha et al. 1965). All size groups of native lake trout in Great Bear Lake appeared to have some members that move and some that do not; Johnson (1975) reported that the movement of individuals ranged from 32 km in one year to no measurable movement for 5 to 6 years.

Walch (1980) used radio tags to investigate the movement of 34 lake trout in Twin Lakes, Colorado. He reported that average movement ranged from 1.1 m/min in fall to 1.6 m/min in summer. During the spring, summer, and fall, movements were significantly greater between 0830 and 1130 than between 2230 and 0500. In the fall, afternoon movements were significantly greater than at night. Home ranges were largest during the spring and fall, and most individuals were found within 3 m of the reservoir bottom. Movement was primarily limited to waters less than 12° C.

Age, Growth, and Food

Maximum age and size. The oldest lake trout reported from the United States was a 28-year-old hatchery fish in Minnesota (Carlander 1969). Rawson (1961) reported 36-year-old lake trout from Lac la Ronge, Saskatchewan. Being long lived, lake trout can reach substantial sizes. Capture records for individual lake trout include: (1) 54.4 kg from Lake Huron; (2) 46.3 kg from Lake Athabaska, Saskatchewan; (3) 39.9 kg from Lake Michigan; (4) 30.16 kg from Great Slave Lake; (5) 28.5 kg from Lake Superior (the sport-catch record); and (6) 11.34 kg from Twin Lakes, Colorado (Sigler and Miller 1963; Carlander 1969; Everhart and Seaman 1971).

Growth. Lake trout fry are about 23 mm total length at time of hatching (Carlander 1969). Growth rates for lake trout vary by year, season, lake area, strain, age, types and densities of food resources available, stock densities (intraspecific competition), and other factors (Martin and Olver 1980).

Weight of lake trout appears to increase at a rate greater than the cube of the length; length-weight relationships and condition factors are generally similar in both sexes (Carlander 1969). In lakes where alewives are the primary source of food, lake trout growth tends to slow with the onset of sexual maturity (age 5 to 7); where larger forage fish species (e.g., ciscoes and suckers) are the primary food, a slowing of growth is not apparent

(Carlander 1969). Growth tends to be slower in the northern part of the range. Lake trout feeding on fish grow faster than those feeding on plankton (Martin 1966). Griest (1977) noted that lake trout reaching legally catchable size (≥ 381 mm) in Twin Lakes, Colorado were usually 4 years old.

Ages of lake trout at first spawning are related to growth rates. Where growth is slow, maturity may not be achieved until 13 to 17 years; in contrast, when growth is rapid, males can mature at age 5 and females at age 6 to 8 (Carlander 1969). In general, males mature one year earlier than females (Martin and Olver 1980). However, when slower growth is caused by plankton feeding, maturity may be reached at a younger age, for example, at age 4 in some Algonquin Park lakes (Martin 1966).

Food. General surveys of the literature suggest that lake trout will tend to feed primarily on the most abundant food available, whether zooplankton, insects, or fish. Salamanders, shrews, and yellow warblers have been found in the stomachs of lake trout (Carlander 1969). Overall, zooplankton and dipteran larvae and pupae are the principal foods of young lake trout; however, even 38 mm individuals may eat small fish (Carlander 1969). Mysis tends to be an important food source in many lakes for all sizes of lake trout, and growth rates have been increased through the introduction of Mysis relicta (Grabowski and Ahern 1982). Larger lake trout generally feed predominantly on fish (Carlander 1969; Frantz and Cordone 1970; Warner 1972; Johnson 1975; Griest 1977; Martin and Olver 1980). However, in many lakes, lake trout continue to feed on zooplankton throughout their lives (Martin 1966; Carlander 1969; Martin and Olver 1980). In Lake Tahoe, crayfish are an important food for lake trout larger than about 380 mm (Frantz and Cordone 1970; Flint and Goldman 1975).

Feeding generally decreases prior to spawning in the fall. Feeding rates tend to remain low during the winter before increasing to maximum in the spring and summer (Carlander 1969; Frantz and Cordone 1970).

Reproduction

Factors affecting reproduction are the same as those affecting growth (Martin and Olver 1980). After reaching sexual maturity, it is not uncommon for lake trout to spawn intermittently (Carlander 1969; Martin and Olver 1980). For example, in Great Bear Lake, lake trout may spawn every third year; whereas in Great Slave Lake spawning may occur every year. Reasons for the intermittent spawning behavior appear to include latitude, temperature, and photoperiod (Martin and Olver 1980).

Spawning periods vary among lake trout population stocks: for example, (1) September to November in the Great Lakes; (2) September to December in New York; (3) October to December in New Hampshire; (4) mid-October to mid-November in Maine; (5) mid-September in Great Slave Lake; and (6) mid-August to early September in Great Bear Lake (Carlander 1969; Johnson 1975). Stocks from different sources may spawn during different periods when planted into the same lake (Griest 1977).

Several reports suggest that declining water temperatures and photoperiod coupled with strong on-shore winds are necessary factors triggering spawning (Carlander 1969; Martin and Olver 1980). Most spawning activity occurs between dusk and 2300 (Carlander 1969; Martin and Olver 1980). Lake trout usually spawn in lakes, rarely in streams.

The average number of eggs in a female lake trout range from 628 to 1710 per kilogram of total weight (Carlander 1969). Average size female lake trout from Jackson Lake, Wyoming, yield as many as 6000 eggs (Baxter and Simon 1970). Hatching time for eggs increases with decreasing temperatures (Embrey 1934; Carlander 1969): 50 days at 10° C; 100 to 117 at 5° C; and 141 to 156 at 2 to 2.5° C.

Specific Habitat Requirements

Lakes containing populations of lake trout have been described as "larger, of higher altitude, deeper, clearer, colder, better oxygenated, more acidic, softer, and lower in total alkalinity, buffering capacity, total dissolved solids and morphoedaphic index values" than waters without lake trout (Martin and Olver 1980). In an analysis of 2500 Ontario lakes, lake trout lakes were distinguished by: (1) mean depths > 6 m; (2) total dissolved solids concentrations < 50 mg/l; (3) average hypolimnetic oxygen concentrations ≥ 6 mg/l; and (4) metric morphoedaphic index values (Ryder 1965) < 6 (Johnson et al. 1977).

Temperature is a critical factor influencing lake trout. During field studies, lake trout are rarely found at water temperatures > 10 to 12° C (e.g., Rawson 1961; Eschmeyer 1964; Johnson 1975; Walch 1980). Carlander (1969) suggested that lake trout can be found at water temperatures ranging between 4 to 18° C during the summer.

Laboratory studies suggest that 12° C is the preferred temperature of adult lake trout, but this preferendum may range between 8 and 15° C (Johnson 1975). For yearling lake trout the preferred temperature was reported as 11.7° C (McCauley and Tait 1970). The preferred temperature for fingerling lake trout was reported as 10.8° C; while that for fry ranged between 9.0 and 10.0° C (Peterson et al. 1979). The upper lethal temperature for adult lake trout was reported to be 23.5° C (Gibson and Fry 1954). Temperature appears to be a stimulus for spawning, though the critical temperature varies between 4.5 and 14° C for different lakes (Rawson 1961; Carlander 1969; DeRoche 1969; Johnson 1975).

For the most part, spawning is limited to lakes, but some spawning has been reported to occur in tributary streams of Lake Superior and in rivers in tundra areas of Canada and Alaska (Loftus 1957; McPhail and Lindsey 1970; Lawrie and Rahrer 1973). In general the beds of the spawning rivers consist of larger boulders intermixed with coarse gravel.

Spawning depths in lakes have been reported to range from 15 cm to over 55 m (Merriman 1939; Carlander 1969; DeRoche 1969; Johnson 1975). For successful spawning in reservoirs it is important that spawning depths remain below drawn-down levels to prevent exposure of eggs (Bergersen and Maiolie 1981).

Lake trout may spawn on a variety of surfaces: clay bottoms in southern Lake Michigan; silt, hardpan clay, marl, and gravel in Green Lake, Wisconsin; and stone bottoms in Lake Superior (Carlander 1969). For most inland lakes spawning occurs primarily on rubble 2.5 cm or larger in diameter (Merriman 1939; Hacker 1956; DeRoche 1969; Martin and Olver 1980). DeRoche (1969) observed lake trout selecting angular rubble over smooth rubble as spawning substrate. Martin (1956) reported spawning areas of lakes in Algonquin Park, Ontario to be free of mud, sand, and detritus. Lake trout have been observed cleaning debris, silt, algae, and slime out of spawning areas prior to spawning (Merriman 1939; DeRoche and Bond 1955; Carlander 1969). Spawning areas for lake trout can be improved by dumping angular rock into lakes (e.g., Hacker 1956).

Dissolved oxygen concentrations of < 6 mg/l have been reported to adversely affect lake trout embryo development and survival at 7 and 10° C (Carlson and Siefert 1974). Garside (1959) suggested that, if oxygen concentrations decrease below 40% of the air saturation at 3° C (about 5.5 mg/l oxygen), lake trout embryos will be subjected to stress and deformities will result. These reports agree with the observation that lake trout tended to be absent from Ontario lakes when average hypolimnetic oxygen concentrations were < 6 mg/l (Johnson et al. 1977).

HABITAT SUITABILITY INDEX (HSI) MODELS

Model Applicability

Geographical area. The models provided are applicable throughout North America except for the Great Lakes. The standard of comparison for each individual variable suitability index is the optimal value of the variable that occurs anywhere within this region. Therefore, the models will never provide an HSI of 1.0 when optimal temperatures, forage fish species, or satisfactory spawning substrate are lacking.

Lake types. The models were developed for oligotrophic (mean metric MEI ≤ 6 , see Ryder 1965) lakes and reservoirs, having mean depths greater than 6 m. The Great Lakes are excluded.

Model Descriptions


Models were developed for (1) stocked lake trout populations, (2) naturally reproducing lake trout populations in lakes, and (3) naturally reproducing lake trout populations in reservoirs. The models were based on different combinations of five habitat variables.

Suitability Index (SI) Graphs for Model Variables

This section contains suitability index graphs for the five variables described below. These graphs are followed by equations for combining selected variable indices, using the component approach, to provide HSI's for stocked lake trout populations in lakes and reservoirs, reproducing populations in lakes, and reproducing populations in reservoirs.


Variable

V₁ Mean maximum temperature in hypolimnion or deeper half of water body.




V₂ Minimum oxygen concentration in hypolimnion or deeper.

- A. O₂ min < 6 mg/l
- B. O₂ ≥ 6 mg/l or ≤ 8 mg/l
- C. O₂ > 8 mg/l




V₃ Availability of forage fish species.


- A. Absent
- B. Present



- V₄ Type and size of available spawning substrate.
- A. Patches of cobble (2 to 30 cm diameter) on lake bottom between 0.5 and 50 m deep.
 - B. Other substrate available that offers eggs and larvae protection from predation.
 - C. Substrate lacking protective cover for eggs and larvae.



- V₅ Annual reservoir drawdown during incubation.
- A. ≤ 0.15 m
 - B. > 0.15 m, but drawdown exposes $< 10\%$ of cobble substrate available for spawning.
 - C. Drawdown exposes 10 to 90% of cobble substrate available for spawning.
 - D. Drawdown exposes 100% of cobble substrate.



Development of Suitability Index Graphs: Rationale and Assumptions

The preceding suitability index graphs should be regarded as tentative and open to modification. The prospective user should understand that the graphs reflect the authors' selective integration of information from the literature, personal experience, and reviewers' comments. The following discussion documents the rationale for constructing the graphs. While there is information on preferred and unsuitable conditions for lake trout, little information exists on intermediate conditions. The models are offered as a starting point, with the hope that refinements will be made as additional information becomes available, including results from model testing.

Mean maximum temperature in hypolimnion (V_1). Lake trout prefer water of 10 to 12° C (Rawson 1961; Eschmeyer 1964; Johnson 1975; Walch 1980), but are abundant in lakes where the water temperature is between 7 and 15° C (Everhart and Seaman 1971; Johnson 1975). Carlander's (1969) review suggested that lake trout are found between 4 and 18° C during the summer. The fish can survive temperatures up to 23.5° C (Gibson and Fry 1954). It is assumed that a maximum summer water temperature of 7 to 15° C is optimum. Above 15° C productivity drops until the upper maximum lethal temperature of 23.5° C is reached. Below 7° C productivity drops due to reduced metabolic rates and depressed community productivity.

Minimum oxygen concentration in hypolimnion (V_2). Dissolved oxygen concentrations of less than 6 mg/l have been shown to depress embryo survival (Garside 1959; Carlson and Siefert 1974) and to be associated with the absence of lake trout in Ontario lakes (Johnson et al. 1977). Therefore, it is assumed that a dissolved oxygen level of 6 mg/l in the hypolimnion of a lake is the absolute minimum required for lake trout survival, but populations subjected to 6 mg/l for extended periods during the growing season would be severely stressed, likely resulting in extensive population mortality and probable population failure.

Presence of forage fish (V_3). Lake trout feed heavily on zooplankton throughout their lives in lakes with little or no fish forage (Martin 1966; Carlander 1969; Martin and Olver 1980). When forage fish are present, lake trout feed predominantly on them (Carlander 1969; Frantz and Cordone 1970; Warner 1972; Johnson 1975; Griest 1977). Lake trout that feed on fish grow faster than those feeding on plankton (Carlander 1969). It is assumed that the productivity of a lake trout population is increased when forage fish are available.

Spawning substrate (V_4). Lake trout can spawn over a variety of substrates (Carlander 1969), but they prefer rubble 2.5 cm or larger in diameter (Merriman 1939; Hacker 1966; DeRoche 1969; Martin and Olver 1980). Spawning areas tend to be free of silt or detritus (Martin 1956). Lake trout have been observed to clean spawning areas of silt and debris prior to spawning (Merriman 1939; DeRoche and Bond 1955). It is assumed that clean rubble (2.5 to 30 cm diameter) is the optimum spawning substrate. Other substrates can produce young if suitable protection from predation is available. Substrates of silt, clay, or marl are generally unsuitable for successful reproduction.

Reservoir drawdown during incubation (V_5). Spawning depths range from 15 cm to > 55 m (Merriman 1939; Carlander 1969; DeRoche 1969; Johnson 1975). Exposure of eggs during incubation prevent their survival (Bergersen and Maiolie 1981). It is assumed that if lake trout spawn in a reservoir they will be impacted by the degree to which the reservoir level fluctuates over this spawning substrate. Where ice is very thick, the values of the suitability index may have to be adjusted to account for habitat exclusion due to ice formation.

HSI Model for Stocked Lake Trout Populations

This model consists of three variables: mean maximum water temperature; minimum oxygen concentration; and presence of forage fish.

$$HSI = V_1 \times V_2 \times V_3$$

HSI Model for Reproducing Lake Trout Populations in Lakes

This model consists of four variables: mean maximum water temperature; minimum oxygen concentration; presence of forage fish; and available spawning substrate.

$$HSI = V_1 \times V_2 \times V_3 \times V_4$$

HSI Model for Reproducing Lake Trout Populations in Reservoirs

This model consists of five variables: mean maximum water temperature; minimum oxygen concentration; presence of forage fish; available spawning substrate; and relative degree of reservoir drawdown.

$$HSI = V_1 \times V_2 \times V_3 \times V_4 \times V_5$$

Interpreting Model Outputs

These lake trout HSI models have not been verified as having a direct relationship to carrying capacity. Although relationships clearly exist between carrying capacities or productivities of lake trout and temperatures and forage fish in the habitat area, insufficient information exists to define these relationships. Therefore, SI's determined for V_1 and V_3 may reflect relative, rather than absolute, differences for populations.

Criteria used for V_4 reflect a "best guess" synthesis of available information. They may be inappropriate for some lakes or reservoirs. Additional data are necessary to refine this predictor. The importance of V_5 can be reduced by the use of deep spawning strains of lake trout when stocking reservoirs.

We believe the models should be able to predict the presence or absence of lake trout as a result of variables V_1 , V_2 , and V_5 . These three variables involve maximum temperature, minimum dissolved oxygen, and maximum exposure of spawning areas that can be encountered by a lake trout population.

ADDITIONAL HABITAT MODELS

Aggus and Bivin (1982) used angler harvest as an indicator of habitat suitability and developed a regression equation relating harvest to reservoir habitat variables for six impoundments in the conterminous United States:

$$\text{Harvest of lake trout} = 1.826 - 0.8714 \log_{10} (\text{outlet depth})$$

Units are kg/ha (harvest) and midline depth of principal outlet in feet below listed surface elevation of reservoir. The authors discuss procedures for converting measured or predicted harvest values to HSI's and present cautionary material concerning model use and interpretation.

REFERENCES

- Aggus, L. R., and W. M. Bivin. 1982. Habitat suitability index models: Regression models based on harvest of coolwater and coldwater fishes in reservoirs. U.S. Fish Wildl. Serv. FWS/OBS-82/10.25. 38 pp.
- Baxter, G. T., and J. R. Simon. 1970. Wyoming fishes. Wyoming Game Fish Dept. Bull. 4, Cheyenne. 168 pp.
- Bergersen, E. P., and M. Maiolie. 1981. Colorado Cooperative Fishery Research Unit studies at Twin Lakes, Colorado: 1980 report of findings. Bur. Rec. Rep. REC-ERC-828:1-66.
- Carlander, K. D. 1969. Handbook of freshwater fishery biology. Iowa State Univ. Press, Ames. 752 pp.
- Carlson, A. R., and R. E. Siefert. 1974. Effects of reduced oxygen on the embryo and larva of lake trout (*Salvelinus namaycush*) and largemouth bass (*Micropterus salmoides*). J. Fish. Res. Board Can. 38:1393-1396.
- Dahlberg, M. D. 1981. Nearshore spatial distribution of fishes in gill net samples, Cayuga Lake, New York. J. Great Lakes Res. 7:7-14.
- DeRoche, S. E. 1969. Observations on the spawning habits and early life of lake trout. Prog. Fish Cult. 31(2):109-113.
- DeRoche, S. E., and L. H. Bond. 1955. The lake trout of Cold Stream Pond Infield, Maine. Trans. Am. Fish. Soc. 85:257-270.
- Embody, G. C. 1934. Relation of temperature to the incubation periods of eggs of four species of trout. Trans. Am. Fish. Soc. 64:281-292.
- Eschmeyer, P. H., R. Daly, and L. F. Erkkila. 1953. The movement of tagged lake trout in Lake Superior, 1950 to 1952. Trans. Am. Fish. Soc. 82:68-77.

- Everhart, W. H., and W. R. Seaman. 1971. Fishes of Colorado. Colorado Game, Fish, and Parks Div., Denver. 75 pp.
- Flint, R. W., and C. R. Goldman. 1975. The effects of a benthic grazer on the primary productivity of the littoral zone of Lake Tahoe. *Limnol. Oceanogr.* 20:935-944.
- Frantz, T. C., and A. J. Cordone. 1970. Food of lake trout in Lake Tahoe. *California Game Fish* 56:21-35.
- Garside, E. T. 1959. Some effects of oxygen in relation to temperature on the development of lake trout embryos. *Can. J. Zool.* 37(5):689-698.
- Gibson, E. S., and F. E. J. Fry. 1954. The performance of the lake trout, Salvelinus namaycush at various levels of temperature and oxygen pressure. *Can. J. Zool.* 32:252-260.
- Grabowski, J. J., and J. Ahern. 1982. Evaluation of limnological parameters as related to the success of Mysis relicta introductions. *Wyoming Water Resour. Res. Inst. Rep.* 85:1-164.
- Griest, J. R. 1977. The lake trout of Twin Lakes, Colorado. *Bur. Rec. Rep.* REC-ERC-77-4:1-29.
- Hacker, V. A. 1956. Biology and management of lake trout in Green Lake, Wisconsin. *Trans. Am. Fish. Soc.* 86:71-83.
- Johnson, L. 1975. Distribution of fish species in Great Bear Lake, Northwest Territories, with reference to zooplankton, benthic invertebrates, and environmental conditions. *J. Fish. Res. Board Can.* 32:1989-2004.
- Johnson, M. G., J. H. Leach, C. K. Minns, and C. H. Olver. 1977. Limnological characteristics of Ontario lakes in relation to associations of walleye (Stizostedion vitreum vitreum), northern pike (Esox lucius), and small-mouth bass (Micropterus dolomieu). *J. Fish Res. Board Can.* 34:1592-1601.
- Khan, N. Y., and S. U. Qadri. 1971. Intraspecies variations and postglacial distribution of lake char (Salvelinus namaycush). *J. Fish. Res. Board Can.* 28:465-476.
- Lawrie, A. H., and J. F. Rahrer. 1973. Lake Superior, a case history of the lake and its fisheries. *Great Lakes Fish. Comm. Tech. Rep.* 19:1-69.
- Loftus, K. H. 1957. Studies on river spawning populations of lake trout in eastern Lake Superior. *Trans. Am. Fish. Soc.* 87:259-277.
- McCauley, R. W., and J. S. Tait. 1970. Preferred temperature of yearling lake trout, Salvelinus namaycush. *J. Fish Res. Board Can.* 27:1729-1733.
- McPhail, J. D., and C. C. Lindsey. 1970. Freshwater fishes of northwestern Canada and Alaska. *Fish. Res. Board Can. Bull.* 173. 381 pp.

- Martin, N. V. 1956. Reproduction of lake trout in Algonquin Park, Ontario. Trans. Am. Fish. Soc. 95:415-422.
- _____. 1966. The significance of food habits in the biology, exploitation, and management of Algonquin Park, Ontario, lake trout. Trans. Am. Soc. 95:415-422.
- Martin, N. V., and C. H. Olver. 1980. The lake charr, Salvelinus namaycush. Pages 205-277 in E. K. Balon (ed.). Charrs - salmonid fishes of the genus Salvelinus. Junk Publ., The Hague, Netherlands.
- Merriman, D. 1939. Squam Lake trout. Boston Soc. Nat. Hist. Bull. 75:3-10.
- Morton, W. M. 1980. Charr or Char: a history of the English name for members of the salmonid genus Salvelinus. Pages 4-6 in E. K. Balon (ed.). Charrs - salmonid fishes of the genus Salvelinus. Junk Publ., The Hague, Netherlands.
- Peterson, R. H., A. M. Sutterlin, and J. L. Metcalfe. 1979. Temperature preferences of several species of Salmo and Salvelinus and some of their hybrids. J. Fish. Res. Board Can. 36:1137-1140.
- Pycha, R. L., W. R. Dryer, and G. R. King. 1965. Movements of hatchery-reared lake trout in southern Lake Superior, 1955-62. J. Fish. Res. Board Can. 24:281-298.
- Rawson, D. S. 1961. The lake trout of Lac la Ronge, Saskatchewan. J. Fish. Res. Board Can. 18:423-462.
- Robins, C. R., R. M. Bailey, C. E. Bond, J. R. Booker, E. A. Lochner, R. N. Lea., and W. B. Scott. 1980. A list of the common and scientific names of fishes from the United States and Canada. Fourth edition. Am. Fish. Soc. Spec. Publ. 12. 174 pp.
- Ryder, R. A. 1965. A method for estimating the potential fish production of north-temperate lakes. Trans. Am. Fish. Soc. 94:214-215.
- Sigler, W. F., and R. R. Miller. 1963. Fishes of Utah. Utah Dept. Fish Game, Salt Lake City. 203 pp.
- Walch, L. A. 1980. Movements of lake trout in Twin Lakes, Colorado, in relation to the Mount Elbert Pumped-Storage Powerplant. Water Power Resour. Serv. Rep. REC-ERC-79-17:1-76.
- Warner, K. 1972. Further studies of fish predation on salmon stocked in Maine lakes. Prog. Fish Cult. 34:217-221.

50272-101		1. REPORT NO. FWS/OBS-82/10.84		2.		3. Recipient's Accession No.	
REPORT DOCUMENTATION PAGE						5. Report Date September 1984	
4. Title and Subtitle Habitat Suitability Index Models: Lake Trout (Exclusive of the Great Lakes)						6.	
7. Author(s) Michael Marcus, Wayne Hubert, and Stanley Anderson						9. Performing Organization Rept. No.	
9. Performing Organization Name and Address Western Aquatic, Inc. P. O. Box 546 Laramie, WY 82070		Wyoming Cooperative Fishery and Wildlife Research Unit University of Wyoming Laramie, WY 82071				10. Project/Task/Work Unit No.	
						11. Contract(C) or Grant(G) No. (C) (G)	
12. Sponsoring Organization Name and Address Western Energy and Land Use Team Division of Biological Services Research and Development Fish and Wildlife Service U.S. Department of the Interior Washington, DC 20240						13. Type of Report & Period Covered	
15. Supplementary Notes						14.	

18. Abstract (Limit: 200 words)

The Habitat Suitability Index (HSI) models for Lake Trout (Salvelinus namaycush) presented in this publication aid in identifying important habitat variables. Facts, ideas, and concepts obtained from the research literature and expert reviews are synthesized into subjective HSI models, which are scaled to produce an index between 0 (unsuitable habitat) and 1 (optimal habitat).

The models are hypotheses of species-habitat relationships, and model users should recognize that the degree of verocity of the HSI model, SI graphs, and assumptions will vary according to geographical area and the extent of the data base for individual variables.

17. Document Analysis a. Descriptors


Fishes
Habitability
Mathematical models

b. Identifiers/Open-Ended Terms

Lake trout
Salvelinus namaycush
Habitat suitability

c. COSATI Field/Group

18. Availability Statement Release unlimited	19. Security Class (This Report) Unclassified	21. No. of Pages 12 pp
	20. Security Class (This Page) Unclassified	22. Price



REGION 1

Regional Director
 U.S. Fish and Wildlife Service
 Lloyd Five Hundred Building, Suite 1692
 500 N.E. Multnomah Street
 Portland, Oregon 97232

REGION 2

Regional Director
 U.S. Fish and Wildlife Service
 P.O. Box 1306
 Albuquerque, New Mexico 87103

REGION 3

Regional Director
 U.S. Fish and Wildlife Service
 Federal Building, Fort Snelling
 Twin Cities, Minnesota 55111

REGION 4

Regional Director
 U.S. Fish and Wildlife Service
 Richard B. Russell Building
 75 Spring Street, S.W.
 Atlanta, Georgia 30303

REGION 5

Regional Director
 U.S. Fish and Wildlife Service
 One Gateway Center
 Newton Corner, Massachusetts 02158

REGION 6

Regional Director
 U.S. Fish and Wildlife Service
 P.O. Box 25486
 Denver Federal Center
 Denver, Colorado 80225

REGION 7

Regional Director
 U.S. Fish and Wildlife Service
 1011 E. Tudor Road
 Anchorage, Alaska 99503



DEPARTMENT OF THE INTERIOR U.S. FISH AND WILDLIFE SERVICE



As the Nation's principal conservation agency, the Department of the Interior has responsibility for most of our nationally owned public lands and natural resources. This includes fostering the wisest use of our land and water resources, protecting our fish and wildlife, preserving the environmental and cultural values of our national parks and historical places, and providing for the enjoyment of life through outdoor recreation. The Department assesses our energy and mineral resources and works to assure that their development is in the best interests of all our people. The Department also has a major responsibility for American Indian reservation communities and for people who live in island territories under U.S. administration.